

daily use of a fluoride preventive treatment” (select one of the following: “rinse” or “gel”) “and a fluoride toothpaste can help reduce the incidence of dental cavities.”

(2) *For dentifrice products containing 1,500 ppm theoretical total fluorine.* “Adults and children over 6 years of age may wish to use this extra-strength fluoride dentifrice if they reside in a nonfluoridated area or if they have a greater tendency to develop cavities.”

[60 FR 52507, Oct. 6, 1995; 60 FR 57927, Nov. 24, 1995; 61 FR 51187, Oct. 7, 1996; 64 FR 13296, Mar. 17, 1999]

**§ 355.55 Principal display panel of all fluoride rinse drug products.**

In addition to the statement of identity required in § 355.50, the following statement shall be prominently placed on the principal display panel: “IMPORTANT: Read directions for proper use.”

**§ 355.60 Professional labeling.**

(a) The labeling for anticaries fluoride treatment rinses identified in § 355.10(a)(3) and (c)(3) that are specially formulated so they may be swallowed (fluoride supplements) and are provided to health professionals (but not to the general public) may contain the following additional dosage information: Children 3 to under 14 years of age: As a supplement in areas where the water supply is nonfluoridated (less than 0.3 parts per million (ppm)), clean the teeth with a toothpaste and rinse with 5 milliliters (mL) of 0.02 percent or 10 mL of 0.01 percent fluoride ion rinse daily, then swallow. When the water supply contains 0.3 to 0.7 ppm fluoride ion, reduce the dose to 2.5 mL of 0.02 percent or 5 mL of 0.01 percent fluoride ion rinse daily.

(b) The labeling for products marketed to health to health professionals in package sizes larger than those specified in § 355.20 shall include the statements: “For Professional Office Use Only” and “This product is not intended for home or unsupervised consumer use.”

**Subpart D—Testing Procedures**

**§ 355.70 Testing procedures for fluoride dentifrice drug products.**

(a) A fluoride dentifrice drug product shall meet the biological test requirements for animal caries reduction and one of the following tests: Enamel solubility reduction or fluoride enamel uptake. The testing procedures for these biological tests are labeled *Biological Testing Procedures for Fluoride Dentifrices*; these testing procedures are on file under Docket No. 80N-0042 in the Division of Dockets Management (HFA-305), Food and Drug Administration, 5630 Fishers Lane, rm. 1061, Rockville, MD 20852, and are available on request to that office.

(b) The United States Pharmacopeia fluoride dentifrice reference standards along with reference standard stability profiles (total fluoride, available fluoride ion, pH, and specific gravity) required to be used in the biological tests are available to any purchaser upon written request to the United States Pharmacopeial Convention, Inc., 1260 Twinbrook Parkway, Rockville, MD 20852.

(c) Alternative testing procedures may be used. Any proposed modification or alternative testing procedures shall be submitted as a petition in accord with § 10.30 of this chapter. The petition should contain data to support the modification or data demonstrating that an alternative testing procedure provides results of equivalent accuracy. All information submitted will be subjected to the disclosure rules in part 20 of this chapter.

[60 FR 52507, Oct. 6, 1995, as amended at 68 FR 24879, May 9, 2003]

**PART 357—MISCELLANEOUS INTERNAL DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE**

**Subpart A [Reserved]**

**Subpart B—Anthelmintic Drug Products**

Sec.

357.101 Scope.

357.103 Definition.

357.110 Anthelmintic active ingredient.